

The effect of subfilter-scale physics on regularization models

Jonathan Pietarila Graham¹, Darryl D. Holm², Pablo Mininni^{3,4},
and Annick Pouquet³

October 12, 2010

¹Max-Planck-Institut für Sonnensystemforschung, Katlenburg-Lindau, Germany ²Department of Mathematics, Imperial College London, London, UK ³National Center for Atmospheric Research,¹ Boulder, Colorado, USA ⁴Departamento de Física, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, Buenos Aires, Argentina

Abstract

The subfilter-scale (SFS) physics of regularization models are investigated to understand the regularizations' performance as SFS models. Suppression of spectrally local SFS interactions and conservation of small-scale circulation in the Lagrangian-averaged Navier-Stokes α -model (LANS- α) is found to lead to the formation of rigid bodies. These contaminate the superfilter-scale energy spectrum with a scaling that approaches k^{+1} as the SFS spectra is resolved. The Clark- α and Leray- α models, truncations of LANS- α , do not conserve small-scale circulation and do not develop rigid bodies. LANS- α , however, is closest to Navier-Stokes in intermittency properties. All three models are found to be stable at high Reynolds number. Differences between L^2 and H^1 norm models are clarified. For magnetohydrodynamics (MHD), the presence of the Lorentz force as a source (or sink) for circulation and as a facilitator of both spectrally nonlocal large to small scale interactions as well as local SFS interactions prevents the formation of rigid bodies in Lagrangian-averaged MHD (LAMHD- α). LAMHD- α performs well as a predictor of superfilter-scale energy spectra and of intermittent current sheets at high Reynolds numbers. It may prove generally applicable as a MHD-LES.

keywords: LES, Subgrid-scale processes, alpha models, MHD, intermittency

¹The National Center for Atmospheric Research is sponsored by the National Science Foundation

1 Introduction

Computing solutions to the Navier-Stokes equations at realistic values of the Reynolds number ($Re \equiv UL_0/\nu$, with characteristic velocity, U , and length-scale, L_0 and viscosity, ν) for most geophysical, astrophysical, and many engineering applications is technologically infeasible. This is because the range of dynamically important length (and time) scales is quite large: from the large scale, L_0 , defined by the problem down to the scales of viscous dissipation, $l_\nu \sim L_0 Re^{-3/4}$. One approach is to simply cut off the smallest scales to arrive at a problem small enough for modern computational limits. A low-bandpass filter, $L : f \rightarrow \bar{f}$, replaces the velocity, $\mathbf{v}$, and pressure, P , with smoother, resolvable fields, $\bar{\mathbf{v}}, \bar{P}$. Their time evolution is governed by the filtered Navier-Stokes equations,

$$\begin{aligned} \partial_t \bar{\mathbf{v}} + \bar{\boldsymbol{\omega}} \times \bar{\mathbf{v}} &= -\nabla \bar{P} + \nu \nabla^2 \bar{\mathbf{v}} - \nabla \cdot \bar{\boldsymbol{\tau}} \\ \nabla \cdot \bar{\mathbf{v}} &= 0, \end{aligned} \quad (1)$$

where $\boldsymbol{\omega} = \nabla \times \mathbf{v}$ is the vorticity and $\bar{\boldsymbol{\tau}}$ is the Reynolds subfilter-scale (SFS) stress tensor,

$$\bar{\tau}_{ij} = \overline{v_i v_j} - \bar{v}_i \bar{v}_j. \quad (2)$$

The modeling of the unresolved stress, Eq. (2), is the main challenge of conducting such large eddy simulations (LES, see [42] for a recent review).

Regularization modeling (of the SFS stress tensor) for Navier-Stokes [10, 29, 14, 17, 8, 18, 33, 40, 50], magnetohydrodynamics (MHD) [28], Boussinesq convection [53], and inviscid cases [37] promises several advantages. For Navier-Stokes, only weak, possibly non-unique solutions have been rigorously proven to exist, and this can impact the possibility of achieving a direct numerical solution (DNS), e.g., with Fourier methods [26]. This is because to prove, even in the linear case, the convergence of a numerical solver for a partial differential equation (PDE), one assumes the PDE is well-posed (i.e., the existence of unique solutions that depend continuously on the data) [38]. A regularization has strong, unique solutions (it is well-posed) and the concerns of [26] do not prevent us from achieving a DNS of the model equations, even with Fourier methods. It is worth noting that numerical convergence of the DNS of the regularization model equations implies a grid-independent model of the Navier-Stokes equations. Additionally, the dissipative term will be unmodified (unlike many LES); the Reynolds number remains well defined (e.g., $\boldsymbol{\omega} \times \bar{\mathbf{v}}/\nu \nabla^2 \bar{\mathbf{v}} \sim UL_0/\nu \equiv Re$). This is opposed to the usual approach of modeling the behavior of the flow in the limit of very large Re . Thus, the models may be more applicable to intermittent phenomena where the length of the inertial range can be important [23]. Since the models do not introduce the effect of the small scales in an *ad hoc* fashion but rather preserve the mathematical properties

of the underlying equations, their application can further our understanding of turbulence and turbulence modeling. The methods are also more easily generalized to other problems (e.g., coupling to a magnetic field).

In this paper we address two separate questions. One is the question of the practical applicability of regularization models as SFS models. When addressing this question, the filter width, α , will be placed in the inertial range and the grid spacing, Δ , will be just small enough to achieve a DNS of the regularization. Our aim is then to determine how well the model’s DNS reproduces a “DNS” of Navier-Stokes compared *at scales larger than* α . Our second question is “How do the models work?” To answer this question, we take an approach that is the antithesis of a practical LES: we choose the filter width, α , to be a large fraction of the computation domain (and, thus, a large multiple of Δ). Such a calculation is not a LES—there is no superfilter-scale inertial range to compare with Navier-Stokes. Instead, we make a new type of study to understand the new SFS physics introduced by the regularization. The differences in physics between the SFS model and Navier-Stokes (or MHD) is what allows the model to reduce computational cost when it is employed as a SFS model. Understanding of how the models work (or fail) can guide the development of new models as we will show.

2 Navier-Stokes

2.1 LANS- α and rigid body formation

The first model we consider is the Lagrangian-averaged Navier-Stokes (LANS) α -model [10, 29]. It is derived by Lagrangian averaging fluid motions followed by application of Taylor’s frozen-in turbulence approximation as the model’s one and only closure: fluctuations about the Lagrangian mean smaller than α are swept along by the large-scale flow and are not allowed to interact with one another [28]. The model is attractive as it retains the Hamiltonian structure of Navier-Stokes, preserves Kelvin’s theorem (conserves small-scale circulation in the absence of dissipation), and conserves both total energy and helicity (the correlation between the velocity, $\mathbf{v}$, and its curl, the vorticity $\boldsymbol{\omega} = \nabla \times \mathbf{v}$) [29]. These properties are conserved in the H_α^1 norm instead of the usual L^2 norm. This is essential when interpreting results of the model as, for example, quantities involving the square velocity, $|\mathbf{v}|^2$, must now be replaced with the dot product $\mathbf{v} \cdot \bar{\mathbf{v}}$ where $\bar{\mathbf{v}}$ is the filtered velocity. Physically, due to the frozen-in approximation, the model retains spectrally nonlocal interactions (important at finite Re [43, 2]) between the superfilter and subfilter scales while the flux of energy in subfilter scales is reduced by the limit on local small-scale to small-scale

interactions [14, 22].

The LANS- α model is given by,

$$\begin{aligned}\partial_t \mathbf{v} + \boldsymbol{\omega} \times \bar{\mathbf{v}} &= -\nabla \pi + \nu \nabla^2 \mathbf{v} \\ \nabla \cdot \mathbf{v} &= \nabla \cdot \bar{\mathbf{v}} = 0.\end{aligned}\tag{3}$$

From the identity, $\boldsymbol{\omega} \times \bar{\mathbf{v}} = \bar{\mathbf{v}} \cdot \nabla \mathbf{v} + (\nabla \bar{\mathbf{v}})^T \cdot \mathbf{v} - \nabla(\bar{\mathbf{v}} \cdot \mathbf{v})$, we see that it differs from Navier-Stokes both in advection by the smoothed velocity and the addition of a second nonlinear term associated with the conservation of circulation. Traditionally, LANS- α is used with an inverse Helmholtz operator as the filter: $\bar{v}_i = (1 - \alpha^2 \partial_{jj})^{-1} v_i$. In this case, LANS- α can be written as a LES, Eq. (1), with

$$\bar{\tau}_{ij}^\alpha = (1 - \alpha^2 \partial_{jj})^{-1} \alpha^2 (\partial_m \bar{v}_i \partial_m \bar{v}_j + \partial_m \bar{v}_i \partial_j \bar{v}_m - \partial_i \bar{v}_m \partial_j \bar{v}_m).\tag{4}$$

The model allows for a reduction in resolution without changing (or supplementing) the dissipative term by instead altering the SFS scaling properties. Near wavenumber, $k = 2\pi/\alpha$, the H_α^1 energy spectrum is predicted to transition from k^β with $\beta = -5/3$ at larger scales to $\beta = -1$ at smaller scales [14]. Consequently, dissipation ($\Omega_{(\alpha)}(k) = k^2 E_{(\alpha)}(k)$) goes as k^1 instead of $k^{1/3}$ and the same amount of power is dissipated in fewer scales. The change in spectral scaling also predicts a resolution requirement in degrees of freedom, dof , for LANS [14],

$$dof_\alpha \sim \alpha^{-1} Re^{3/2},\tag{5}$$

which has been confirmed in numerical experiments [22]. Once dof_α has been resolved, further resolution yields no change in the numerical solution: LANS- α is a grid-independent SFS model. When compared with the dof for Navier-Stokes,

$$dof_{NS} \sim Re^{9/4},\tag{6}$$

we see that LANS- α should improve as a SFS model for larger Re . This was an encouraging prediction as LANS- α compared well with dynamic eddy viscosity [46] and dynamic mixed (similarity) eddy viscosity [18] at moderate Re .

We have, however, found that LANS- α develops a problem at large Re : it develops a positive-exponent power-law bump in its small-scale energy spectrum and a contamination of superfilter-scale spectral properties [22]. To investigate the SFS physics responsible for this, we employ (as the antithesis of a LES) a filter $1/3$ the size of our 256^3 computational cube in a pseudo-spectral calculation [20, 21] with a Taylor-Green (TG) forcing [51] and $Re \approx 8000$. As shown in Fig. 1, the observed scaling law is k^{+1} . This was shown to be associated with the formation in the flow of passively swept regions, called *rigid bodies* [22]. These form as a consequence




Figure 1: Energy spectrum for LANS- α (solid line) with filter scale $\alpha = 2\pi/3$ (vertical dashed line). The SFS power law is well approximated by k^{+1} . Insert shows thresholded cubed velocity increment $|\delta v_{\parallel}(2\pi/10)|^3 < 10^{-2}$ in black. These regions do not contribute to the turbulent cascade of energy to smaller scales and are identified with rigid bodies. A spectrum of only the white regions (dashed line) is consistent with the predicted k^{-1} scaling outside rigid bodies.

of disallowing sub- α -scale fluctuations to interact with each other in the closure approximation. A rigid body cannot support longitudinal velocity increments: $\delta \bar{v}_{\parallel}(l) \equiv [\bar{\mathbf{v}}(\mathbf{x}) - \bar{\mathbf{v}}(\mathbf{x} + \mathbf{l})] \cdot \mathbf{l}/l = 0$. This dimensionally predicts a scaling relation, $\delta \bar{v} \sim l^0$, and, with $v \sim (1 + \alpha^2 k^2) \bar{v} \sim \alpha^2 k^2 \bar{v}$ for $l \ll \alpha$, an energy spectrum of

$$E_{\alpha}(k) \sim \bar{v} v k^{-1} \sim k^1 \quad (7)$$

which is compatible with the observed SFS energy spectrum. Inside rigid bodies there can be no turbulent cascade of energy to smaller scales (since there are no internal degrees of freedom). From the Kármán-Howarth theorem, we should then expect to be able to detect rigid bodies by visualizing the cubed velocity increments (which are proportional to the energy flux). The regions which correspond to negligible flux are shown as black in the inset of Fig. 1. Filtering these regions out, allows us to obtain a (convolved) energy spectra for the remaining white portion of the flow. This spectrum is shown as a dashed line in Fig. 1 and has a negative spectral slope close to the predicted k^{-1} spectrum. The resulting picture of the model's behavior is to produce two spatially separate scalings. The white portions of the flow possess the predicted LANS- α scaling and are responsible for the observation of the predicted dof_{α} . The black portions are rigid bodies whose k^1 energy spectrum dominates over k^{-1} for large k and are responsible for the observed spectral contamination. Because of this spectral contamination, there is no need for further tests of LANS- α against newer models such as the Variational Multiscale Method [30]. Note, however, that suitable spectra can be obtained with a modified viscous length scale (LANS- $\alpha\beta$ [33]).

2.2 The influence of circulation on rigid bodies

The formation of rigid bodies in LANS- α limits the reduction of numerical *dof* saved compared to Navier-Stokes to a factor of 1/12 regardless of Re [22]. It is desirable, then, to alter the model in such a way to prevent the formation of rigid bodies. Truncation of the SFS stress tensor, Eq. (4), to the first term results in the Clark- α model [8],

$$\begin{aligned} \partial_t \mathbf{v} + (1 - \frac{1}{2} \alpha^2 \nabla^2)(\bar{\omega} \times \bar{\mathbf{v}}) - \frac{1}{2} \alpha^2 [(\nabla^2 \bar{\omega}) \times \bar{\mathbf{v}} + \bar{\omega} \times (\nabla^2 \bar{\mathbf{v}})] = \\ -\nabla \mathcal{P} + \nu \nabla^2 \mathbf{v} \\ \nabla \cdot \mathbf{v} = \nabla \cdot \bar{\mathbf{v}} = 0, \end{aligned} \quad (8)$$

and to the first two terms results in the Leray- α model [17, 18],

$$\begin{aligned} \partial_t \mathbf{v} + \bar{\mathbf{v}} \cdot \nabla \mathbf{v} = -\nabla p + \nu \nabla^2 \mathbf{v} \\ \nabla \cdot \mathbf{v} = \nabla \cdot \bar{\mathbf{v}} = 0. \end{aligned} \quad (9)$$

Both these models are regularizations and conserve the total energy of the flow [8, 17, 18]. They do not, however, conserve the helicity nor the small-scale circulation. Considering the rotational properties of a rigid body (in the absence of viscous friction), these models' circulation properties may be incompatible with rigid body formation. Indeed, while LANS- α exhibits a positive-exponent power law in this case ($\alpha = 2\pi/13$, $Re \approx 3300$, TG forcing), both Clark- α and Leray- α are free from this signature of rigid body formation (Fig. 2a). However, LANS- α 's intermittency properties are more similar to Navier-Stokes than the other two models (Fig. 2b).

2.3 Computability and interpretation of H_α^1 norm regularizations

It has been reported recently that Clark- α and LANS- α have poor SFS model performance and, in fact, have instabilities in simulations with periodic boundary conditions [9]. This result highlights the importance of correctly interpreting H_α^1 norm regularizations. To illustrate this, we compute 256^3 DNS and 64^3 models runs for their case 3a: decaying Taylor-Green vortex with $\nu = 2.5 \cdot 10^{-3}$ (time step, $dt = 3 \cdot 10^{-3}$ for all simulations). In Figs. 3 and 4, we display results for the models employing a filter width $\alpha = 2\pi/16$ (results with $\alpha = 2\pi/24$ are closer to that of the DNS). Fig. 3a shows the time evolution of the total kinetic energy in the super-filter scales,

$$tke \equiv \int_0^{k_\alpha} E_{(\alpha)}(k), \quad (10)$$

where $k_\alpha = 2\pi/\alpha$. Incorrectly applying the L^2 norm, $E = |\mathbf{v}|^2/2$, to Clark- α and LANS- α leads to the interpretation that they deviate significantly from the DNS due to slower energy decay. Correctly applying the H_α^1 norm, $E_\alpha = \bar{\mathbf{v}} \cdot \mathbf{v}/2$, shows that the models are, in fact, doing quite well as can be seen in the plot of the dissipation rate of energy, ε (Fig. 3b). The qualitative properties of Lagrangian models is that they have the same invariants as the primitive equations, albeit in a different norm. Since it is these invariants which may very well influence the dynamics, as for example shown in [12] where Kolmogorov spectra are present at intermediate times at large scale in ideal (Euler) three-dimensional fluid flows, one must compare the behavior of one set of invariants to the other set in the Lagrangian modeling formulation. Similarly, the L^2 norm spectra for Clark- α and LANS- α in Fig. 4a appear under-dissipative at high wave numbers, while the correct H_α^1 norm spectra are closer to the DNS. We also conduct Clark- α and LANS- α computations for case 3b of [9]: $\nu = 1/3000$, $\alpha = 2\pi/32$ and 384^3 grid points (Fig. 4b). We find no signs of instability. The particular numerical expression of the models we used




Figure 2: (a) Compensated energy spectra ($2\pi/\alpha$, vertical dashed line) for Navier-Stokes (solid black), LANS- α (blue dash-dotted), Clark- α (green dashed), and Leray- α (red dotted). (b) Normalized structure function scaling exponent ξ_p/ξ_3 versus order p . Clark- α is the best approximation for the superfilter-scale spectrum, whereas high-order intermittency properties are best reproduced by LANS- α [24].




Figure 3: Case3a, $Re \approx 400$. Line styles are as in Fig. 2a for correct norms with the addition of L^2 norm energy for Clark- α (pink dash-triple-dotted) and for LANS- α (cyan long-dashed). (a) Filtered kinetic energy, tke . (b) Dissipation, $\varepsilon = dE/dt$, versus time, t .

is given in Eqs. (3) and (8). These differ from Eqs. (8) and (9) in [9] and it is known that discretized pseudospectral operators depend on their algorithmic form (e.g., $\nabla\psi^2 \neq 2\psi\nabla\psi$ [13, 7]). This is likely the source of their observed instability; however a recently discovered deficiency in high-order low-storage Runge-Kutta schemes [6] may also impact numerical implementation of the models. For this reason, our calculations are made with second-order Runge-Kutta in time. It is worth pointing out that Clark- α is nearly identical to the Rational Large Eddy Simulation (RLES) model [15] which has been shown to be unstable for non-periodic boundaries [5, 32]. Both our work and that of [9] have employed periodic boundary conditions only, but extra precautions must be taken for stability in the non-periodic case [5, 32].

3 MHD: circulation and outlook for LES

In MHD, the circulation properties are quite different since small-scale circulation is broken by the Lorentz force $\mathbf{j} \times \mathbf{b}$, with $\mathbf{j} = \nabla \times \mathbf{b}$ the current, $\mathbf{b}$ being the induction. This force acts as source (sink) of circulation, Γ , as opposed to the insufficient modeling of Γ in the Leray- α and Clark- α models. This can be seen in Kelvin's theorem,

$$\frac{d}{dt}\Gamma = \frac{d}{dt} \oint_{\mathcal{C}} \mathbf{v} \cdot d\mathbf{r} = \oint_{\mathcal{C}} \mathbf{j} \times \mathbf{b} \cdot d\mathbf{r}. \quad (11)$$

This may prevent the formation of rigid bodies even while conserving all the correct physical properties of the flow.

The LES equations for MHD are given by

$$\begin{aligned} \partial_t \bar{\mathbf{v}} + \bar{\boldsymbol{\omega}} \times \bar{\mathbf{v}} &= \bar{\mathbf{j}} \times \bar{\mathbf{b}} - \nabla \bar{\Pi} + \nu \nabla^2 \bar{\mathbf{v}} - \nabla \cdot \bar{\boldsymbol{\tau}} \\ \partial_t \bar{\mathbf{b}} &= \nabla \times (\bar{\mathbf{v}} \times \bar{\mathbf{b}}) + \eta \nabla^2 \bar{\mathbf{b}} - \nabla \cdot \bar{\boldsymbol{\tau}}^b, \end{aligned} \quad (12)$$

where η is the magnetic diffusivity, $\Pi = P + |\mathbf{b}|^2/2$ the modified pressure, $\bar{\boldsymbol{\tau}}$ is the Reynolds SFS stress tensor,

$$\bar{\tau}_{ij} = \overline{v_i v_j} - \bar{v}_i \bar{v}_j - (\overline{b_i b_j} - \bar{b}_i \bar{b}_j), \quad (13)$$

and $\bar{\boldsymbol{\tau}}^b$ is the electromotive-force SFS stress tensor,

$$\bar{\tau}_{ij}^b = \overline{b_i v_j} - \bar{b}_i \bar{v}_j - (\overline{v_i b_j} - \bar{v}_i \bar{b}_j). \quad (14)$$

Note that the extension of eddy viscosity to eddy resistivity employs the usual form for $\bar{\boldsymbol{\tau}}$ involving only the filtered velocity while the expression for $\bar{\boldsymbol{\tau}}^b$ similarly only involves the filtered magnetic field [52]. Meanwhile, Eqs. (13) and (14) make it explicitly clear that interactions between the two fields *at subfilter scales* must be taken into account.




Figure 4: (a) Case3a, $Re \approx 400$: energy spectra after peak of dissipation, $t \in [4.5, 5.5]$. Line styles are as in Fig. 3. Vertical dashed line is $2\pi/\alpha$. (b) Case3b, $Re \approx 3000$: dissipation, ε , versus time, t for Clark- α (light green) and for LANS- α (dark blue).

Another problem with extending eddy-viscosity concepts to MHD is that they can be related to a known power law of the energy spectrum [11]. This is inappropriate for MHD as neither kinetic nor magnetic energy is a conserved quantity and the general expression of the energy spectrum is not known at this time [31, 35, 19, 41, 39]. Additionally, MHD has been shown to have nonlocal interactions between large and small (superfilter and subfilter) scales [1] (e.g., in the Batchelor viscous-inductive regime, $\nu \gg \eta$, where energy is transferred directly from viscous-scale, l_ν , eddies to small-scale magnetic field, $l \ll l_\nu$ [3]). This complex interaction is a challenge in general for MHD-LES, but may be an advantage for the Lagrangian-averaged approach as energy exchanges with sub- α scales may disrupt rigid body formation. Some limited case MHD LES include the cross-helicity model [48] assuming alignment between the fields and the low magnetic Re LES [34, 49]. Extensions of spectral models to MHD based on two-point closure formulations of the dynamical equations proposed recently look promising in the analysis of turbulent flows and of the dynamo mechanism [4]. Approximate deconvolution models for MHD [36] also appear promising, but have yet to be tested on non-laminar flows. However, there are many regimes of MHD dependent on the ratios between the various conserved quantities and ν/η : there has yet to be demonstrated a generally applicable LES for MHD.

3.1 LAMHD- α and absence of rigid bodies

The Lagrangian-averaged MHD α -model (LAMHD- α) [28, 27, 47] is given by, where the velocity is filtered as before and $\bar{\mathbf{b}} = (1 - \alpha_M^2 \nabla^2)^{-1} \mathbf{b}$:

$$\begin{aligned} \partial_t \mathbf{v} + \boldsymbol{\omega} \times \bar{\mathbf{v}} &= \mathbf{j} \times \bar{\mathbf{b}} - \nabla \pi + \nu \nabla^2 \mathbf{v} \\ \partial_t \bar{\mathbf{b}} &= \nabla \times (\bar{\mathbf{v}} \times \bar{\mathbf{b}}) + \eta \nabla^2 \bar{\mathbf{b}} \\ \nabla \cdot \mathbf{v} &= \nabla \cdot \bar{\mathbf{v}} = \nabla \cdot \mathbf{b} = \nabla \cdot \bar{\mathbf{b}} = 0. \end{aligned} \tag{15}$$

LAMHD- α may be written as a MHD-LES, Eqs. (12), for the case $\alpha_M = \alpha$ [23, 25], which we study here. The model preserves the ideal quadratic invariants of MHD (in the H_α^1 norm) as well as Alfvén's theorem for frozen-in field lines [28]. Physically, it supports Alfvén waves at all scales while slowing and hyper-diffusively damping waves with wavelengths, λ , smaller than α [25]. In examinations of its SFS physical properties LAMHD- α (dashed lines) displays neither positive-exponent power-law scaling nor superfilter-scale spectral contamination (see Fig. 5). Under similar conditions LANS- α (not shown) displays these signs of rigid body formation. Further examinations with larger filters and higher Re were unable to unravel any sign that rigid bodies form for LAMHD- α [25].




Figure 5: Kinetic (a) and magnetic (b) energy spectra ($2\pi/\alpha$ vertical dashed line). The largest scales are affected by differences from the MHD DNS (solid lines) in initial conditions. LAMHD- α (dashed lines) exhibits neither the positive power-law nor the superfilter-scale spectral contamination associated with high Re LANS- α .

3.2 LAMHD- α as a SFS model

Since LAMHD- α did not display any signs of rigid body spectral contamination, we test it as a SFS model for large kinetic and magnetic Reynolds numbers, ≈ 3300 . A DNS of MHD is computed at a resolution of 1024^3 . The initial conditions for $\mathbf{v}$ and $\mathbf{b}$ are a superposition of ABC modes [16] with random phases and wavenumbers $k \in [1, 4]$. No external forcing is applied and the total energy is allowed to freely decay. LAMHD- α is computed for identical conditions at a resolution of $N^3 = 168^3$ with the same Re and a filter size $\alpha = 6\Delta = 2\pi/28$. As a base-level comparison we also compute an under-resolved (or no-model) solution of the MHD equations at $N^3 = 168^3$. Time evolution of the total energies and enstrophy are shown in Fig. 6. In comparison with under-resolving MHD, LAMHD- α shows errors of approximately the same magnitude in these global quantities. Comparisons of energy spectra (Fig. 7), however, show an improvement in predictive quality for LAMHD- α , especially for the magnetic energy spectrum. As turbulence develops, energy begins to pile up at small-scales and deplete at intermediate scales for 168^3 MHD. LAMHD- α improves the prediction of superfilter-scale spectra compared to no SFS model.

Cross-sections of $|\mathbf{j}|^2$, shown in Fig. 8 at $t = 8.4$ indicate that LAMHD- α finds sharper and better defined, more intermittent current sheets than the under-resolved run compared to the DNS.

4 Summary

Incompressible LANS- α , while it performed well at moderate Reynolds number is limited as a high Re SFS model. Due to its strong suppression of spectrally local interactions at subfilter-scales, and consistent with its conservation of small-scale circulation, LANS- α develops rigid bodies which contaminate the superfilter-scale energy spectrum. In contrast, Clark- α and Leray- α , neither of which conserve small-scale circulation do not develop energy-spectrum contamination from rigid bodies. LANS- α , however, best matches the intermittency properties of Navier-Stokes fluid turbulence.

In MHD, a mechanism for local small-scale transfer is the interaction of small-scale Alfvén waves. As LAMHD- α supports Alfvén waves at all scales while slowing and hyperdiffusively damping those with wavelength $\lambda < \alpha$, it more gently suppresses SFS local interactions than LANS- α . This together with the greater nonlocality in MHD and the Lorentz-force source of small-scale circulation, inhibits the formation of rigid bodies in LAMHD- α . It appears to retain the good intermittency properties of LANS- α without its poor spectral properties. For this reason, we find LAMHD- α to be a viable model at high Re in 3D. As LAMHD- α has been




Figure 6: Time evolution of total (solid), magnetic (dashed), and kinetic (dotted) energies (a) and total dissipation, ϵ , (b) for 1024³ DNS (black/solid), 168³ LAMHD- α (blue/dash-dotted), and 168³ no-model (red/dotted). LAMHD- α provides no improvement in prediction of global quantities over an under-resolved DNS.




Figure 7: Compensated kinetic (a) and magnetic (b) energy spectra averaged over $t \in [8, 8.8]$. Line styles are as in Fig. 6. ($2\pi/\alpha$ vertical dashed line). Energy piles up at small scales in the no-model approach (under-resolved DNS) and LAMHD- α is seen to provide improved predictions of the superfilter-scale spectra, especially for the magnetic field.




Figure 8: Cross sections of square current, j^2 , at $t = 8.4$ for no-model 168^3 (left), 168^3 LAMHD- α (center), and 1024^3 DNS (right) at full resolution. LAMHD- α provides a much better capturing of the intermittent current sheets than the under-resolved solution.

previously found to reproduce the difficult to model properties of MHD at high Re in 2D [45] and moderate Re in 3D [44], we believe it will prove to be a generally applicable MHD LES, in many instances in geophysics and astrophysics where magnetic fields are known to be important dynamically.

References

- [1] A. Alexakis, P. D. Mininni, and A. Pouquet. Shell-to-shell energy transfer in magnetohydrodynamics. I. Steady state turbulence. *Phys. Rev. E*, 72(4):046301–+, October 2005.
- [2] H. Aluie and G. L. Eyink. Localness of energy cascade in hydrodynamic turbulence. II. Sharp spectral filter. *Physics of Fluids*, 21(11):115108–+, November 2009.
- [3] H. Aluie and G. L. Eyink. Scale-Locality of Magnetohydrodynamic Turbulence. *ArXiv e-prints*, December 2009.
- [4] J. Baerenzung, H. Politano, Y. Ponty, and A. Pouquet. Spectral modeling of magnetohydrodynamic turbulent flows. *Phys. Rev. E*, 78(2):026310–+, August 2008.
- [5] Luigi C. Berselli and Carlo R. Grisanti. On the consistency of the Rational Large Eddy Simulation model. *Computing and Visualization in Science*, 6(2-3):75–82, March 2004.
- [6] M. E. Brachet, P. D. Mininni, D. L. Rosenberg, and A. Pouquet. High-order low-storage explicit Runge-Kutta schemes for equations with quadratic nonlinearities. *ArXiv e-prints*, August 2008.

- [7] Claudio Canuto, M. Yousuff Hussaini, Alfio Quarteroni, and Thomas A. Zang. *Spectral Methods in Fluid Dynamics*. Springer-Verlag, New York, 1988.
- [8] C. Cao, D. D. Holm, and E. S. Titi. On the Clark α model of turbulence: global regularity and long-time dynamics. *Journal of Turbulence*, 6:N20, 2005.
- [9] A. Chandy and S. Frankel. Regularization-based sub-grid scale (SGS) models for large eddy simulations (LES) of high-Re decaying isotropic turbulence. *Journal of Turbulence*, 10:25–+, 2009.
- [10] S. Chen, C. Foias, D. D. Holm, E. Olson, E. S. Titi, and S. Wynne. Camassa-Holm Equations as a Closure Model for Turbulent Channel and Pipe Flow. *Physical Review Letters*, 81:5338–5341, December 1998.
- [11] J.-P. Chollet and M. Lesieur. Parameterization of Small Scales of Three-Dimensional Isotropic Turbulence Utilizing Spectral Closures. *Journal of Atmospheric Sciences*, 38:2747–2757, December 1981.
- [12] C. Cichowlas, P. Bonaïti, F. Debbasch, and M. Brachet. Effective Dissipation and Turbulence in Spectrally Truncated Euler Flows. *Physical Review Letters*, 95(26):264502–+, December 2005.
- [13] J. P. Dahlburg, D. Montgomery, G. D. Doolen, and W. H. Matthaeus. Large-scale disruptions in a current-carrying magnetofluid. *Journal of Plasma Physics*, 35:1–42, February 1986.
- [14] C. Foias, D. D. Holm, and E. S. Titi. The Navier-Stokes-alpha model of fluid turbulence. *Physica D Nonlinear Phenomena*, 152-153:505–519, May 2001.
- [15] G. P. Galdi and W. J. Layton. Approximation of the larger eddies in fluid motions. II: A model for space-filtered flow. *Math. Models Methods Appl. Sci.*, 10:343–350, 2000.
- [16] D. Galloway and U. Frisch. Dynamo action in a family of flows with chaotic streamlines. *Geophysical and Astrophysical Fluid Dynamics*, 36:53–83, 1986.
- [17] B. J. Geurts and D. D. Holm. Regularization modeling for large-eddy simulation. *Physics of Fluids*, 15:L13–L16, January 2003.
- [18] Bernard J. Geurts and Darryl D. Holm. Leray and LANS- α modelling of turbulent mixing. *Journal of Turbulence*, 7(10):1–33, 2006.

- [19] P. Goldreich and S. Sridhar. Toward a theory of interstellar turbulence. 2: Strong Alfvénic turbulence. *Astrophys. Journal*, 438:763–775, January 1995.
- [20] D. O. Gómez, P. D. Mininni, and P. Dmitruk. MHD simulations and astrophysical applications. *Advances in Space Research*, 35:899–907, 2005.
- [21] D. O. Gómez, P. D. Mininni, and P. Dmitruk. Parallel Simulations in Turbulent MHD. *Physica Scripta Volume T*, 116:123–127, January 2005.
- [22] J. Pietarila Graham, D. Holm, P. Mininni, and A. Pouquet. Highly turbulent solutions of the Lagrangian-averaged Navier-Stokes alpha model and their large-eddy-simulation potential. *Phys. Rev. E*, 76:056310–+, 2007.
- [23] J. Pietarila Graham, D. D. Holm, P. Mininni, and A. Pouquet. Inertial range scaling, Kármán-Howarth theorem, and intermittency for forced and decaying Lagrangian averaged magnetohydrodynamic equations in two dimensions. *Physics of Fluids*, 18:045106, April 2006.
- [24] J. Pietarila Graham, D. D. Holm, P. D. Mininni, and A. Pouquet. Three regularization models of the Navier-Stokes equations. *Physics of Fluids*, 20(3):035107–+, March 2008.
- [25] J. Pietarila Graham, P. D. Mininni, and A. Pouquet. Lagrangian-averaged model for magnetohydrodynamic turbulence and the absence of bottlenecks. *Phys. Rev. E*, 80(1):016313–+, July 2009.
- [26] J.-L. Guermond. On the use of the notion of suitable weak solutions in CFD. *International Journal for Numerical Methods in Fluids*, 57:1153–1170, July 2008.
- [27] D. D. Holm. Averaged Lagrangians and the mean effects of fluctuations in ideal fluid dynamics. *Physica D Nonlinear Phenomena*, 170:253–286, September 2002.
- [28] D. D. Holm. Lagrangian averages, averaged Lagrangians, and the mean effects of fluctuations in fluid dynamics. *Chaos*, 12:518–530, 2002.
- [29] D. D. Holm, J. E. Marsden, and T. S. Ratiu. The Euler-Poincaré Equations and Semidirect Products with Applications to Continuum Theories. *Adv. in Math.*, 137:1–81, 1998.

- [30] T. J. R. Hughes, L. Mazzei, A. A. Oberai, and A. A. Wray. The multiscale formulation of large eddy simulation: Decay of homogeneous isotropic turbulence. *Physics of Fluids*, 13:505–512, February 2001.
- [31] P. S. Iroshnikov. Turbulence of a Conducting Fluid in a Strong Magnetic Field. *Soviet Astronomy*, 7:566–+, February 1964.
- [32] V John. An assessment of two models for the subgrid scale tensor in the rational LES model. *Journal of Computational and Applied Mathematics*, 173(1):57–80, January 2005.
- [33] T.-Y. Kim, M. Cassiani, J. D. Albertson, J. E. Dolbow, E. Fried, and M. E. Gurtin. Impact of the inherent separation of scales in the Navier-Stokes- $\alpha\beta$ equations. *Phys. Rev. E*, 79(4):045307–+, April 2009.
- [34] B. Knaepen and P. Moin. Large-eddy simulation of conductive flows at low magnetic Reynolds number. *Physics of Fluids*, 16:1255–+, May 2004.
- [35] R. H. Kraichnan. Inertial-range spectrum of hydromagnetic turbulence. *Physics of Fluids*, 8:1385–1387, 1965.
- [36] A. Labovschii and C. Trenchea. Approximate deconvolution models for magnetohydrodynamics. Technical report, University of Pittsburgh, 2010.
- [37] A. Larios and E. S. Titi. On the Higher-Order Global Regularity of the Inviscid Voigt-Regularization of Three-Dimensional Hydrodynamic Models. *ArXiv e-prints*, October 2009.
- [38] P. D. Lax and R. D. Richtmyer. Survey of the stability of linear finite difference equations. *Communications on Pure and Applied Mathematics*, 9:267293, May 1956.
- [39] E. Lee, M. E. Brachet, A. Pouquet, P. D. Mininni, and D. Rosenberg. Lack of universality in decaying magnetohydrodynamic turbulence. *Phys. Rev. E*, 81(1):016318–+, January 2010.
- [40] B. Levant, F. Ramos, and E. S. Titi. On the Statistical Properties of the 3D Incompressible Navier-Stokes-Voigt Model. *ArXiv e-prints*, January 2009.
- [41] J. Mason, F. Cattaneo, and S. Boldyrev. Numerical measurements of the spectrum in magnetohydrodynamic turbulence. *Phys. Rev. E*, 77(3):036403–+, March 2008.

- [42] C. Meneveau and J. Katz. Scale-Invariance and Turbulence Models for Large-Eddy Simulation. *Annual Review of Fluid Mechanics*, 32:1–32, 2000.
- [43] P. D. Mininni, A. Alexakis, and A. Pouquet. Nonlocal interactions in hydrodynamic turbulence at high Reynolds numbers: The slow emergence of scaling laws. *Phys. Rev. E*, 77(3):036306–+, March 2008.
- [44] P. D. Mininni, D. C. Montgomery, and A. Pouquet. Numerical solutions of the three-dimensional magnetohydrodynamic α model. *Phys. Rev. E*, 71(4):046304–+, April 2005.
- [45] P. D. Mininni, D. C. Montgomery, and A. G. Pouquet. A numerical study of the alpha model for two-dimensional magnetohydrodynamic turbulent flows. *Physics of Fluids*, 17(3):035112–+, March 2005.
- [46] K. Mohseni, B. Kosović, S. Shkoller, and J. E. Marsden. Numerical simulations of the Lagrangian averaged Navier-Stokes equations for homogeneous isotropic turbulence. *Physics of Fluids*, 15:524–544, February 2003.
- [47] D. C. Montgomery and A. Pouquet. An alternative interpretation for the Holm “alpha model”. *Physics of Fluids*, 14(9):3365–3366, 2002.
- [48] W.-C. Müller and D. Carati. Dynamic gradient-diffusion subgrid models for incompressible magnetohydrodynamic turbulence. *Physics of Plasmas*, 9:824–834, March 2002.
- [49] Y. Ponty, H. Politano, and J.-F. Pinton. Simulation of Induction at Low Magnetic Prandtl Number. *Physical Review Letters*, 92(14):144503–+, April 2004.
- [50] F. Ramos and E. S. Titi. Invariant measures for the 3D Navier-Stokes-Voigt equations and their Navier-Stokes limit. *ArXiv e-prints*, October 2009.
- [51] G. I. Taylor and A. E. Green. Mechanism of the Production of Small Eddies from Large Ones. *Proceedings of the Royal Society of London*, A158:499, 1937.
- [52] M. L. Theobald, P. A. Fox, and S. Sofia. A subgrid-scale resistivity for magnetohydrodynamics. *Physics of Plasmas*, 1:3016–3032, September 1994.
- [53] M. van Reeuwijk, H. J. J. Jonker, and K. Hanjalić. Leray- α simulations of wall-bounded turbulent flows. *International Journal of Heat and Fluid Flow*, 30:1044, 2009.